Waters™



Drugs of Abuse - LC/UV

Waters Corporation



For forensic toxicology use only.

This is an Application Brief and does not contain a detailed Experimental section.

Abstract


This application brief demonstrates analysis of drugs of abuse - LC/UV.

Introduction

The compounds used in this study are -


Compounds	USP Tailing
3,4-Methylenedioxyamphetamine (MDA)	1.08
3,4-Methylenedioxymethamphetamine (MDMA)	1.32
Alprazolam	1.18
Flunitrazepam	1.15
Desmethyidiazepam	1.14
Diazepam	1.13

3,4-Methylenedioxymethamphetamine (MDMA)



Diazepam

Alprazolam



3,4-Methylenedioxyamphetamine (MDA)

Flunitrazepam



Desmethyidiazepam

2


Experimental

Drugs of Abuse - LC/UV

Conditions

Column: Xterra RP $_{18}$ 3.0 x 100 mm, 3.5 μm Part number: 186000420 Mobile phase A: 5 mM NH₄HCO₃, pH 9.6 Mobile phase B: ACN Flow rate: 0.6 mL/min Isocratic mobile phase composition: 70% A; 30% B Injection volume: 10 μL Sample concentration: 10 ng/μL 40 °C Temperature: Detection: UV @ 210 nm Instrument: Alliance 2690, 2996 PDA

Results and Discussion



Featured Products

· Alliance HPLC System https://www.waters.com/534293

WA20738.046, June 2002



© 2021 Waters Corporation. All Rights Reserved.