


Nota applicativa

Ranitidine in Rat Plasma

Waters Corporation




This is an Application Brief and does not contain a detailed Experimental section.

Abstract

This application brief demonstrates analysis of ranitidine in rat plasma.

Introduction

The compound analyzed in this study is ranitidine.



Ranitidine


Experimental

Conditions

| | |
|-------------------------------------|---|
| Column: | Xterra MS C ₁₈ 2.1 x 30 mm, 3.5 µm |
| Part number: | 186000398 |
| Mobile phase A: | 100 mM NH ₄ COOH |
| Mobile phase B: | ACN |
| Isocratic mobile phase composition: | 30% A; 70% B |
| Flow rate: | 0.2 mL/min |
| Injection volume: | 15 µL |

| | |
|--------------------------|---|
| Detection: | MS ESI+ |
| Instrument: | Alliance 2790, Micromass Quattro Ultima |
| Ion source: | ESI+ |
| Source temperature: | 150 °C |
| Gas cell: | 1.5e-3 mbar, 20eV |
| Desolvation temperature: | 350 °C |
| Cone gas flow: | 150 L/hr |
| Drying gas flow: | 600 L/hr |
| Cone voltage: | 20V |

Oasis® MCX Extraction Method
Oasis® MCX Extraction Plate, 10 mg/96-well
Part Number 186000259




Results and Discussion

CID mass spectra


cid 15
S3 1 (1.005)

Daughters of 315ES+
2.99e8



IS daughter of 271.0
2 88 (1.639) Cm (87:90)

Daughters of 271ES+
6.87e4



Featured Products

Alliance HPLC System <<https://www.waters.com/534293>>

WA20738.096, June 2002

