Waters™

アプリケーションノート

Phenolics in Echinacea, Commercial Products or Whole Leaf

Waters Corporation



Abstract

This application brief highlights the analysis of phenolics using Oasis MAX method.

Introduction

Phenolic compounds in Echinacea, Commercial Products or Whole Leaf are analyzed in this application brief.

CHLOROGENIC ACID

Experimental

Sample Preparation

100 mg sample is extracted with 25 mL of ethanol:water (70:30)

1 mL of the ethanolic extract is diluted 1:3 with water (~ pH 7)

- Recovery, measured with certified standards at 100 ppm in reagent water, was greater than 85% for all compounds
- All analyses gave results within \pm 35% of the expected values with the exception of the liquid supplement (see goldenseal analysis)
- The selective SPE extraction and cleanup procedure provided a convenient analysis of echinacea phenolics in complex matrix such as in herbal tea



HPLC Method

Detection:

Column: Symmetry C_{18} , 4.6 x 250 mm, 5 μ m (p/n: WAT054275) 0.1% phosphoric acid Mobile phase A: Mobile phase B: Acetonitrile **Gradient:** 10% B initial, then linear gradient to 22% B in 13 min, to 40 % B in 40 min Flow rate: 1.5 mL/min Injection volume: $10 \, \mu L$ Temperature: 35 °C


Oasis MAX Methodfor Phenolics in Echinacea

Conditions for Oasis MAX Cartridge, 3 cc/60 mg Part Number 186000367 < https://www.waters.com/nextgen/in/en/shop/sample-preparation--filtration/186000367-oasis-max-3-cc-vac-cartridge-60-mg-sorbent-per-cartridge-30--m-1.html>


UV @ 330 nm

OASIS® MAX METHOD FOR PHENOLICS IN ECHINACEA

Conditions for Oasis® MAX Cartridge, 3 cc/60 mg Part Number 186000367



Results and Discussion



Featured Products

WA31764.121, June 2003

© 2021 Waters Corporation. All Rights Reserved.