# Waters™

# Pharmaceutical Compounds in Plasma

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

#### **Abstract**


This application brief demonstrates analysis of pharmaceutical compounds in plasma.

## Introduction


The compounds used in this study are -

- 1. Caffeine
- 2. Practolol
- 3. N-acetyl procainamide
- 4. Propranolol
- 5. Methoxyverapamil
- 6. Amitriptyline


# Caffeine



N-acetyl procainamide



Methoxyverapamil



### Practolol

Propranolol

Amitriptyline

## Experimental

#### LC Conditions

Column: Xterra MS  $C_{18}$ , 2.1 x 20 mm  $\emph{IS}$ , 3.5  $\mu$ m, (p/n:

186001923)

Mobile phase A: 10 mM NH<sub>4</sub>HCO<sub>3</sub>, pH 10

Mobile phase B: Methanol

Flow rate: 0.4 mL/min

Injection volume: 20  $\mu$ L

Sample concentration: 5 µg/mL

Temperature: Ambient

Detection: MS

Instrument: Waters 277 Sample Manager, Waters 1525

Binary HPLC Pump and Waters Micromass

Quattro Ultima

#### Gradient

Time (min)	Profile	
	% <b>A</b>	%B
0.0	100	0
5.0	5	95

#### **MS Conditions**

ES <sup>+</sup> MRM cone (V):	5.0
Capillary:	3.5
Source temp. (°C):	150
Desolvation temp. (°C):	400
Cone gas flow (L/Hr):	50
Desolvation gas fLow(L/Hr):	550
LM resolution 1 & 2:	13.5
HM resolution 1 & 2:	13.5
lon energy 1:	0.4
lon energy 2:	0.8
Multiplier (V):	650
Sample Preparation	
SPE:	Oasis HLB µElution Plate, (p/n:186001828BA) p/n:186001828BA
Condition:	200 μL Methanol
Equilibrate:	200 μL Water
Load:	150 $\mu$ L Spiked rat plasma with 2% $H_3PO_4$ , diluted with 1:1 Water

Quattro Ultima


Wash:	200 µL 5% Methanol in Water
Elute:	50 μL Methanol
Dilute:	100 µL Water

#### Results and Discussion

Compounds (MRM Transitions, Cone voltage (V), CID (eV)

- 1. Caffeine (195 > 138; 20; 20)
- 2. Practolol (267 > 190.1; 30; 20)
- 3. N-acetyl procainamide (278 > 205.1; 25;20)
- 4. Propranolol (260 > 183; 25;18)
- 5. Methoxyverapamil (485.2 > 165.1; 45; 30)
- 6. Amitriptyline (278 > 233.1; 25; 17)

The top figure is the total ion current, followed by the extracted ion signals for each of the three analytes.



### **Featured Products**

WA31787.19, June 2003

© 2021 Waters Corporation. All Rights Reserved.