Waters™

アプリケーションノート

Antibacterial Drugs in TFA: 900 μg Load

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

Abstract


This application brief highlights the SunFire mass loading 1 2 3 comparison of antibacterial drugs in TFA with other leading HPLC column brands.

Introduction

SunFire mass loading 1 2 3 comparison of antibacterial drugs in TFA with other leading HPLC column brands

Compounds used for this study:

- 1. Oxacillin
- 2. Cloxacillin
- 3. Dicloxacillin



Experimental

Conditions

Column: SunFire C_{18} 4.6 x 150 mm, 5 μm

Part number: 186002559

Mobile phase A: 0.1% TFA in Water

Mobile phase B: 0.1% TFA in ACN

Flow rate: 1.4 mL/min

Injection volume: $10~\mu L$

Sample concentration: 30 mg/mL each, in DMSO

Temperature: Ambient


Detection: UV @ 254 nm

Instrument: AutoPurification System

Gradient

Time (min)	Profile	
	%A	%B
0.0	70	30
10.0	10	90

Results and Discussion



*: $W_{1/2}$ value is calculated based on the width at half-height of the $3^{\rm rd}$ peak.

Featured Products

AutoPurification System https://www.waters.com/10007147

WA41883, May 2005

© 2021 Waters Corporation. All Rights Reserved.