Waters™

Note d'application

Lansoprazole: Isocratic Separation and Degradation by 0.4 N HCl

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

Abstract

This application brief highlights the isocratic separation and degradation of lansoprazole.

Introduction

Lansoprazole is used to treat ulcers, gastroesophageal reflux disease (GERD), and conditions where the stomach produces too much acid.

Experimental

Conditions

Column: SunFire C_{18} 4.6 x 150 mm, 5.0 μ m (p/n:

186002559)

Mobile phase A: 20 mM HCOO-NH₄ +, pH 3.0

Mobile phase B: Acetonitrile

Isocratic: as indicated

Flow rate: 1.4 mL/min

Injection volume: $2 \mu L$

Sample Diluent:	75:25 H ₂ O:ACN
Sample concentration:	350 μg/mL
Temperature:	30 °C
Detection:	UV @ 254 nm
Sampling rate:	10 pts/sec
Time Constant:	0.1
Instrument:	Waters Alliance HT 2795, with 2996
Conditions	
Column:	SunFire C ₁₈ 4.6 x 150 mm, 5.0 µm (p/n: 186002559)
	SunFire C_8 4.6 x 150 mm, 5.0 μ m (p/n: 186002737)
Mobile phase A:	20 mM HCOO-NH ₄ +, pH 3.0
Mobile phase B:	Acetonitrile
Isocratic:	as indicated
Flow rate:	1.4 mL/min
Injection volume:	5 μL
Sample Diluent:	50:50 H ₂ O:ACN
Sample concentration:	2.63 mg/mL

Temperature: 30 °C

Detection: UV @ 254 nm

Sampling rate: 5 pts/sec

Time Constant:

Instrument: Waters Alliance HT 2795, with 2996

Degradation Conditions:


Temperature: ambient


50 mg of Lansoprazole + 5 mL of 0.4N HCl stirred for \sim 30 seconds

Stop reaction by add 0.9 mL of 0.4N NaOH, then dilute with 1.9 mL ACN

Lansoprazole degraded ~ 32%

Results and Discussion





0.00

0.00

2.00

4.00

6.00


8.00

Minutes

10.00

12.00

14.00



Featured Products

Alliance HPLC https://www.waters.com/514248

WA41893, March 2005

© 2021 Waters Corporation. All Rights Reserved.