Waters™

응용 자료

Advil Allergy Sinus Tablet: Oasis MCX

Waters Corporation



Abstract


This application note highlights analysis of advil allergy sinus tablet usig Oasis MCX.


Introduction

This formulation is a good example of a mixture of bases and an acid. The Oasis 2x4 Method was employed and the Oasis MCX plates resulted in the best recoveries for all three analytes.

The compounds used in this study are -


- · Pseudoephedrine
- · Chlorpheniramine
- · Ibuprofen





Pseudoephedrine (Base) MW 165.2 pKa 9.9

Chlorpheniramine (Base) MW 274.8 pKa 9.2



lbuprofen (Acid) MW 206.3 pKa 5.2

Experimental

Test Conditions


Oasis MCX 10-mg 96-well Plates


Condition:	500 μL MeOH	
Equilibrate:	500 μL H ₂ O	
Load:	500 μL sample (250 μL plasma diluted 1:1 with 4% $H_3 PO_4$ in $H_2 O)$	
Wash 1:	500 μL 2% FA	
Elute 1:	2 x 125 µL MeOH (Ibuprofen)	
Elute 2:	$2x125~\mu L$ 5% NH_4OH in MeOH (Bases)	
Options:	1. Dilute Elute 2 with 250 μL 2% FA in water and Elute 1 with 250 μL 100% water and analyze separately.	
	 Combine the two elutions and evaporate/reconstitute. 	
Pseudoephedrine HCI (Base):		
Pseudoephedrine HCl (Base): Chlorpheniramine Maleate (Base):	evaporate/reconstitute.	
	evaporate/reconstitute. 1.5 μg/mL	
Chlorpheniramine Maleate (Base):	evaporate/reconstitute. 1.5 μg/mL 0.1 μg/mL	
Chlorpheniramine Maleate (Base): Ibuprofen (Acid):	evaporate/reconstitute. 1.5 μg/mL 0.1 μg/mL 10 μg/mL	
Chlorpheniramine Maleate (Base): Ibuprofen (Acid): Column:	evaporate/reconstitute. 1.5 μg/mL 0.1 μg/mL 10 μg/mL ACQUITY UPLC BEH C ₁₈ , 2.1 x 50 mm, 1.7 μm	

Oasis MCX 10-mg 96-well Plates


Injection volume:	10.0 μL
Column temp:	4 °C
Sample temp:	10 °C
Instrument:	ACQUITY UPLC with Quattro Premier


SPE Recovery Data: Optimun Sorbent and Protocol





Oasis WAX (Prxotocol 1)





Gradient


Time	Profile		
(min)	%A	%B	
0.0	95	5	
1.0	95	5	
3.0	20	80	
4.0	20	80	
4.5	95	5	
6.0	95	5	

Quattro Premier

ESI ⁺ and ESI ⁻ capillary:	3.0 kV
Source temp:	100 °C
Desolvation temp:	350 °C
Cone gas flow:	0 L/Hr
Desolvation gas flow:	700 L/Hr
Collision cell pressure:	2.59 e ⁻³ mbar

Compound	Precursor ion (<i>m/z</i>)	Product ion (<i>m/z</i>)	Cone voltage (V)	Collision energy (eV)
Ibuprofen (ESI-)	205.1	161.1	20	12
Pseudoephedrine (ESI+)	166.2	117	30	20
Clorpheniramine (ESI+)	275.2	232.2	30	20

Results and Discussion



Featured Products

ACQUITY UPLC System <https://www.waters.com/514207>

WA60091, June 2007

© 2021 Waters Corporation. All Rights Reserved.