Waters™

응용 자료

Analysis of Food Additives and Preservatives using XBridge Phenyl

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

Abstract

This application brief highlights the analysis of food additives and preservatives using XBridge Phenyl columns.

Introduction

Compounds used in this study are: 1. Saccharin 2. p- Hydroxybenzoic Acid 3. Sorbic Acid 4. Methylparaben 5. Dehydroacetic Acid

1. Saccharin

2. p-Hydroxybenzoic Acid

3. Sorbic Acid

4. Methylparaben


5. Dehydroacetic Acid

Experimental

Test Conditions

Columns:	XBridge Phenyl, 4.6 x 100 mm, 3.5 μm p/n: 186003334
Mobile phase A:	20 mM KH ₂ PO ₄ , pH 2.5
Mobile phase B:	ACN
Flow rate:	1.0 mL/min
Isocratic Mobile Phase Composition:	75% A; 25% B
Injection volume:	10 μL
Sample:	Saccharin (100 μ g/mL), P- hydroxybenzoic Acid (10 μ g/mL),
	DehydroAcetic Acid (100 μg/mL),
	Methylparaben (25 μg/mL),
	Sorbic Acid (10 μ g/mL), in KH ₂ PO ₄ /ACN (75/25)
Column temp.:	30 °C
Sampling temp.:	15 °C
Detection:	UV @ 254 nm
Sampling Rate:	5 points/sec
Filter Response:	0.2
Instrument:	Alliance 2695 with 2996 PDA

Results and Discussion



Compounds: 1. Saccharin 2. p- Hydroxybenzoic Acid 3. Sorbic Acid 4. Methylparaben 5. Dehydroacetic Acid.

Featured Products

Alliance HPLC https://www.waters.com/514248

WA60192, June 2007

© 2021 Waters Corporation. All Rights Reserved.