
Waters™

Note d'application

Microcystins in Natural Waters

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

Abstract

Microcystin-LR is a potent mammalian toxin which is known to have been responsible for the deaths of domesticated animals, livestock loss, and the potential presence in potable water supplies.

Introduction

Microcystin-LR is a potent mammalian toxin which is known to have been responsible for the deaths of domesticated animals, livestock loss, and the potential presence in potable water supplies.


Experimental

Pretreatment

- 1. Filter water sample through 0.45 μm membrane filter.
- 2. Add 100 μ L of enkephalin (concentration 10 μ g/L) to 10 mL filtered water sample and mix thoroughly.

SPE Procedure

Oasis® HLB 3 cc/60 mg



LC Conditions

System:	Alliance HPLC 2695
Column:	Symmetry300 C ₁₈ , 3.5 μm, 4.6 x 75 mm
Flow rate:	0.2 mL/min
Mobile phase A:	0.2% formic acid in water
Mobile phase B:	0.2% formic acid in methanol
Injection volume:	10 µL

Column temp.: 30 °C

Gradient

Time (min)	%A	%В
0	45	55
12	10	90
12.5	0	100
15	0	100
15.1	45	55
25	45	55

MS Conditions

MS System: Waters Quattro Ultima Pt

Ionization mode: Positive electrospray (ESI⁺)

Multiple reaction monitoring

Analyte	MRM	MW	[M+H] ⁺	[M+H] ²⁺	Characteristic Ion Fragment
Falsashalia	FF61 . 279.0	EEEC	EEG 1	N.D	278.0
Enkephalin	556.1 → 278.0	555.6	556.1		397.1
MCYST-LR	519.9 → 135.0	994.5	995.7	498.4	135.0
MC131-LR	519.9 → 135.0	994.5	995.7	496.4	861.5
MCYST-RR	498.4 → 135.0	1037.6	1038.4	519.9	135.0
MC131-KK	490.4 → 135.0	1031.0	1036.4		620.0
MCYST-LW	1025.8 → 891.7	1024.5	1025.8	N.D	897.1
IMCT 31-LW	1025.6 → 691.7	1024.5	1025.6		583.2
MCYST-LF	MCYST-LF 986.8 → 852.5 985.5 986	986.8	N.D	852.5	
MCISI-LF	900.0 → 002.0	300.0	900.0	ט.א	544.0

Results and Discussion

Analyte	Concentration (µg/L)	Average Recovery (%)	RSD (%)
	0.10	100.0	6.45
MCYST-RR	0.20	95.2	4.02
	0.40	90.0	4.35
MCYST-LR	0.02	105.0	5.40
	0.05	96.0	4.53
	0.08	93.8	4.22
MCYST-LW	0.40	103.8	5.30
	1.00	102.7	5.87
	1.60	93.8	5.67
MCYST-LF	0.20	103.0	7.03
	0.50	109.8	5.69
	0.80	102.3	4.57

Recovery data for spiked samples at various concentrations.

References

 Determination of Microcystins in Natural Water by Liquid Chromatography Tandem Mass Spectrometry, Chen Qi, Huang Baifen, Zhang Jing, Ren Yiping; Zhejiang Provincial Center for Disease Prevention and Control.

Featured Products

Alliance HPLC System https://www.waters.com/534293

720002595, April 2008

© 2021 Waters Corporation. All Rights Reserved.