
## Waters™

# Multi-Residue Analysis of Pesticides in Teas Using Modified AOAC QuEChERS Method by UPLC-MS/MS

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

#### **Abstract**

This application highlights Multi-Residue Analysis of Pesticides in Tea extracts by UPLC-MS/MS

## Experimental

#### Sample prep procedure

- 1. Tare weigh an empty beaker.
- 2. Weigh out 100 g of tea leaves in the beaker.
- 3. Add in 600 g of hot water at 80-85 °C to the beaker. Brew the tea for 20 minutes.
- 4. Weigh the beaker with water and tea.
- 5. Calculate the weight of water loss due to evaporation. Add water to the beaker to make up for the loss of water.
- 6. Homogenize the sample until it reaches consistent texture.

#### **Extraction Procedure**

- 1. Transfer 15 g of homogenized sample into an empty 50 mL tube.
- 2. Add any internal standards and standard mixture.
- 3. Add 15 mL 1% acetic acid in acetonitrile into the 50 mL DisQuE extraction tube 1.
- 4. Transfer all the powder in the DisQuE extraction tube 1 into the 50 mL containing sample and solvent.
- 5. Shake vigorously for 1 minute and centrifuge > 1500 rcf for 5 minute.
- 6. Transfer 1 mL of the acetonitrile extract into the clean-up tube 2.
- 7. Shake for 30 seconds and centrifuge >1600 rcf for 5 minute.
- 8. Transfer 100  $\mu$ L of final extract into an autosampler vial.

- 9. Add any post-extraction internal standards.
- 10. Dilute as needed with an appropriate buffer or solvent.

#### LC conditions

LC System:

Column:	ACQUITY UPLC BEH C <sub>18</sub> , 2.1 x 100 mm, 1.7 μm
Column Temp:	40 °C

Waters ACQUITY UPLC System

Sample Temp: 4 °C

Flow Rate: 0.3 mL/min.

Mobile Phase A: Water + 0.1% formic acid

Mobile Phase B: Methanol + 0.1% formic acid

Injection Volume: 15  $\mu$ L, Partial loop injection

#### Gradient

Time	Flow	A%	В%
	Rate		
0	0.3	75	25
0.25	0.3	75	25
7.75	0.3	5	100
8.5	0.3	0	100
8.51	0.5	75	25

Time	Flow	A%	В%
	Rate		
10.5	0.5	75	25
11	0.3	75	25

### MS conditions

Instrument: Waters ACQUITY TQ Detector


Ionization: Positive electrospray (ESI+)

Acquisition: Multiple reaction monitoring

(MRM)

## Results and Discussion

## Pesticides Recovery in Teas by UPLC-MS/MS



Pesticides in Teas by UPLC-MS/MS

## **Featured Products**

ACQUITY UPLC System <a href="https://www.waters.com/514207">https://www.waters.com/514207</a>

720003365, March 2010

©2019 Waters Corporation. All Rights Reserved.