Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

Paula Hong, Patricia R. McConville

Waters Corporation

Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

1

This is an Application Brief and does not contain a detailed Experimental
section.

Abstract

This application highlight demonstrates an assay for the analysis of abacavir and related substances that was
successfully transferred from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System.

Introduction

When transferring a method to a different HPLC/UHPLC instrument, the new instrumentation must provide the
same separation and meet all of the system suitability requirements of the original method/instrument. For
gradient separations, the impact of dwell volume can be dramatic and should be considered when transferring a
method across chromatographic instruments.

Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient
SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

Experimental

LC Conditions

Sample: Abacavir and related substances

Sample diluent: Water

Systems: 1. Agilent 1100 Series LC System

                            2. ACQUITY UPLC H-Class System with ACQUITY
                            UPLC PDA Detector and CH-A

Transfer: Gradient delay was entered using SmartStart
                            Technology

Wavelength: 245 nm

Column temp.:   36 °C

Column: CORTECS C_{18}, 2.7 μm, 4.6 x 75 mm

Mobile phase A: 0.1% Trifluoroacetic acid in water

Mobile phase B: 85% Methanol in water

Flow rate: 1.85 mL/min

Injection volume: 5 μL

Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient
SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

3

Gardient:

Time (min)	%A	%B	%C
Initial	95	5	Initial
6.38	70	30	6.38
10.37	10	90	10.37

Results and Discussion

Name	t_R Agilent 1100	t_R ACQUITY UPLC H-Class
1. Descycleopropyl abacavir	4.23	4.35
2. Abacavir	6.34	6.4
3. 1R, 4R Trans abacavir	6.64	6.7
4. O-(4-Chloro-2,5-diaminopyrimidynl)-abacavir	6.93	6.99
5. O-t-Butyl-abacavir	7.70	7.73

Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

4

Agilent 1100 Series LC System

Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

5

ACQUITY UPLC H-Class System

Conclusion

An assay for the analysis of abacavir and related substances was successfully transferred from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System. Using gradient SmartStart Technology in the instrument method, the differences in system volume were factored into the method. No changes to the gradient table were required.

References

1. For complete experimental details, refer to full technical brief 720005252EN at waters.com

Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

Featured Products

- [ACQUITY UPLC H-Class PLUS System <https://www.waters.com/10138533>](https://www.waters.com/10138533)

- [ACQUITY UPLC PDA Detector <https://www.waters.com/514225>](https://www.waters.com/514225)

720005253, December 2014

∧

Method Transfer from an Agilent 1100 Series LC System to an ACQUITY UPLC H-Class System with Gradient SmartStart Technology: Analysis of an Active Pharmaceutical Ingredient and Related Substances[1]

7