Waters™

应用纪要

Isolating Trace Impurities for Structural Elucidation in a Commercial Fungicide Formulation Using Preparative SFC

John McCauley, Marian Twohig

Waters Corporation

Abstract

Trace impurities in synthetic products that interact with human end users or may have an undesirable environmental fate are regulated by various government agencies such as FDA and EPA. As a result, impurity separation and their structural identification are important research functions for many industries, including pharmaceutical, agrochemical, food, and consumer products. Full chemical identification requires structural elucidation of the separated compound using high resolution mass spectrometry (HRMS). However, MS alone is often insufficient to unambiguously identify a compound, especially in the case of isomers. This often necessitates obtaining the isolated pure compounds of interest using purification procedures for study using NMR spectroscopy. In this application note, we describe how a workflow to achieve the full structural elucidation of trace impurities can be implemented using preparative supercritical fluid chromatography to isolate trace impurities.

Benefits

- Supercritical fluid chromatography (SFC) offers high speed and efficiency, as well as the environmentally sustainable benefit of lower solvent consumption.
- This workflow is ideal for industries facing escalating pressure for improved productivity while adhering to strict deadlines and budget constraints.
- Improved enantiomeric and diasteriomeric resolution and shorter analysis times can be obtained using the ACQUITY UPC² System.

Introduction

Trace impurities in synthetic products that interact with human end users or may have an undesirable environmental fate are regulated by various government agencies such as FDA and EPA. As a result, impurity separation and their structural identification are important research functions for many industries, including pharmaceutical, agrochemical, food, and consumer products. Full chemical identification requires structural elucidation of the separated compound using high resolution mass spectrometry (HRMS). However, MS alone is often insufficient to unambiguously identify a compound, especially in the case of isomers. This often necessitates obtaining the isolated pure compounds of interestusing purification procedures for study using NMR spectroscopy.

In this application note, we describe how a workflow to achieve the full structural elucidation of trace impurities can be implemented using preparative supercritical fluid chromatography to isolate trace impurities. A commercial formulation of the fungicide propiconazole was used as an example to demonstrate this workflow.¹ Propiconazole has the structural potential for the existence of several stereoisomers. The propiconazole product used contains isomers (Figure 1) and also related trace impurities at approximately the 1% level (Figure 2). Some propiconazole impurities have previously been structurally identified.² The SFC based workflow described here is generally applicable for impurity isolation and offers many advantages, including high speed and efficiency, fast dry-down, quick turnaround time, as well as the environmentally sustainable benefit of lower solvent consumption.³



Figure 1. Structures of the propiconazole stereoisomers.



Figure 2. Achiral UPC² analysis of the propiconazole in the pesticide formulation sample using an ACQUITY UPC² BEH Column showing two peaks for the propiconazole and two impurity peaks (top trace). Lower trace: chiral analysis using an ACQUITY UPC² Trefoil AMY1 Column showing four peaks for propiconazole and four peaks for the trace impurities.

Experimental

Sample preparation

A representative sample was obtained from a commercial formulation containing 1.55% of propiconazole and its trace impurities, using liquid-liquid extraction with dichloromethane (DCM) and a 5% sodium bicarbonate solution (NaHCO₃).

Analytical and preparative chromatography conditions

(See Table 1)

	Achiral UPC ² Analysis	Chiral UPC ² Analysis	Achiral SFC 80 Prep	Chiral SFC 100 Prep	
Separation mode	Isocratic	Isocratic	Isocratic	Isocratic	
Column	ACQUITY UPC ² BEH 3.0 x 150 mm, 1.7-µm	ACQUITY UPC ² Trefoil AMY1 3.0 x 150 mm, 2.5-µm	Waters Viridis BEH 19 x 150 mm, 5-µm	Chiralpak IC or AD-H 30 x 150 mm, 5-µm	
Co-solvent/Method	Methanol	2-Propanol/Ethanol	Methanol Methanol		
conditions	6% Methanol in CO_2	10% iPrOH/EtOH in CO ₂	5% Methanol in CO ₂	IC: 18% Methanol in CO ₂	
				AD-H: 6% Methanol in CO ₂	
ABPR	120 bar	137 bar	120 bar	120 bar	
Flow rate	1.5 mL/min	2.0 mL/min	70 mL/min	100 mL/min	
UV detection	222 nm	222 nm	222 nm	222 nm	
Column temp.	40 °C	45 °C	40 °C	40 °C	

Table 1. A summary of analytical UPC² and preparative chromatographic conditions. The use of both the SFC 80q Preparative System and the Prep 100q SFC System was based on the dimensions of the columns used in this study.

Mass spectral detection was performed on a Waters SQ Detector 2 in ESI mode scanning from 200 to 700 Da with a capillary voltage of 3 kV, cone voltage of 30 V, source temp. of 150 °C, and a desolvation temp. of 450 °C using an extracted ion at 342 Da, (M+H)⁺.

NMR methods

¹H ¹³C, and 2D NMR spectra were acquired at 400 MHz using a Bruker Avance III spectrometer at the COSMIC Laboratory at Old Dominion University in Norfolk, VA. Assignments were based on chemical shifts, splitting patterns, integration of the peaks, coupling constants, coupling patterns in a 2D COSY spectrum (Correlated Spectroscopy) and one-bond carbon-proton coupling correlations observed in an HSQC spectrum (Heteronuclear Single Quantum Coherence) edited for carbon multiplicities. In a 2D NOESY spectrum (Nuclear Overhauser Effect Spectroscopy), NOE (Nuclear Overhauser Effect) was observed between the protons at carbons 11 and 14 of the triazole and the protons on carbon 6. The same NOE signal was observed in a 1D NOE spectrum with selective irradiation of the triazole protons in impurity peak 1 from the ACQUITY UPC² BEH Column. There were no equivalent NOE signals observed in impurity peak 2.

Results and Discussion

A commercial fungicide formulation (150 mL) containing propiconazole as the active ingredient at a concentration of 1.55% was suspended in 5% NaHCO₃ solution then extracted with DCM three times. The combined extracts were dried over sodium sulphate (Na₂SO₄), filtered, then concentrated. The expected amount (2.3 g) of active ingredient (AI) was contained in 4.5 g of crude extract that contained other inactive ingredients including various surfactants. These crude extracts were then analyzed using UPC². In both achiral and chiral modes (Figure 2) trace impurities at <1% of the amount of the original 1.55% claim were observed.

Methods were then developed to scale separations to preparative chromatography. The achiral prep chromatography was carried out using an SFC 80q Preparative System with a 19 mm I.D. Viridis BEH Column. The analysis showed that the two impurity peaks were obtained in 98% purity (Figure 3).



Figure 3. Preparative collections of trace impurity peaks using a 19 mm I.D. Viridis BEH Column (top

trace). UPC² analysis of the two impurity peak batches resulting from the SFC 80q preparative analysis demonstrate that these collections resulted in >98% purity for each impurity fraction (lower trace).

Sufficient pure material was collected both to enable full structural assignments of the trace impurities as well as to carry out further chiral separations of each peak into their individual enantiomers. MS analysis confirmed that the two isolated impurity peaks were isobaric with each other as well as with the main ingredient suggesting that they were indeed structural isomers.

¹H, ¹³C, 2D, and NOE NMR experiments revealed that these two impurities differed from propiconazole itself by the nitrogen attachment point of the triazole moiety to the methylene group on the dioxolane ring (Figure 4). This result is clearly evident due to symmetry in the NMR. The propiconazole, being attached at the N adjacent to the other N is not symmetric, while in the impurity the N has a C atom adjacent on both sides of the nitrogen, giving symmetry to the triazole moiety and a simpler NMR. The assignment of the cis and trans isomers results from the fact that a strong NOE is seen in both a 2D NOESY NMR experiment as well as in a 1D NOE spectrum between the protons attached to carbons 11 and 14 and those on carbon 6 in the isomer assigned as cis, while this same NOE was not observed in the NMR spectrum of the isomer assigned as trans.



Figure 4. Structures shown for propiconazole (left) and for its impurities (top right, cis isomers, Impurity 1 from Figure 3), and (bottom right, trans isomers, Impurity 2 from Figure 3). Note that the parent propiconazole triazole is connected at a different nitrogen atom.

Having completed the full structural assignments for the two impurity peaks that could be resolved with achiral chromatography, attention was next turned to preparatively separating the enantiomers for both the cis and trans isomers (Figure 5). While all four isomers can be resolved in a single run analytically on a small particle size chiral column, in this case two different column chemistries were required for the separation of each enantiomeric pair upon scale up. The cis isomer separated on the IC column chemistry while the trans isomer required an AD-H column. All four of the possible impurity stereoisomers were then obtained in their pure form.



Figure 5. Preparative collections of the enantiomers of the cis isomer on the Chiralpak IC Column and of the trans isomer on the Chiralpak AD-H Column (left) using the Prep 100q SFC System. UPC² chiral analysis demonstrates that all four of the possible impurity stereoisomers were each preparatively isolated in the enantiopure form with >98% enantiomeric excess (right).

Chiral analysis using UPC² demonstrated that each isomer had an enantiomeric excess >98%. The cis isomer was comprised of the 2R, 4S and the 2S, 4R isomers, while the trans isomers contained the 2R, 4R and the 2S, 4S isomers. Although the assignment of the absolute configurations for each of the stereoisomers was beyond the scope of this study, these isolated impurities do in fact possess the same structures as those found in a previous study.² The capability to have all four of the impurity stereoisomers in hand in their pure form, in sufficient supply, then allows for the unambiguous assignment of the mechanistic potencies and toxicities of each individual impurity species. This capability then enables improved knowledge of the efficacy and safety of the product mixture ingredients.

Conclusion

ACQUITY UPC² analyses of actual product mixtures, using both chiral and achiral column chemistries are useful for both impurity profiling and also for developing separation methods suitable for eventual scale up to preparative separations.

Trace impurities, as in this case <1% of a 1.55% formulation of the active ingredient, can be efficiently isolated in the amounts needed for structural elucidation and other needed studies using preparative SFC.

The isolated impurities can be obtained in their pristine forms, thus enabling the simple application of HRMS, 1D, and 2D NMR spectroscopic studies, allowing for the full structural elucidations of these trace impurities.

Initially isolated impurities can be further separated into their enantiomeric pairs through the implementation of chiral stationary phase preparative SFC with high efficiency.

The assigned structures for all of the isolated impurities were in agreement with information in the prior literature.²

References

- 1. G von Reet, J Heeres, and L Wals Janssen. Ger Pat. 2:551–560, 1976.
- R Glaser, I Adin, D Ovadia, E Mendler, M Drouin. Solid-state structure determination and solutionstate NMR characterization of the (2R,4R)/2S,4S)- and (2R,4S)/(2S,4R)-diastereomers of the agricultural fungicide propiconazole, the (2R,4S)/(2S,4R)-symmetrical triazole constituational isomer, and a ditriazole analogue. *Structural Chemistry*. (6) 3:145–156, 1995.
- 3. R Chen. Mass-directed preparative SFC: An orthogonal toll with reduced liquid solvent usage for high throughput purification. Waters application note no. 720003772en. October 2010.

Acknowledgement

Special thanks for the intellectual contributions of James E. Hall, NMR Spectroscopist, Old Dominion University, Norfolk, VA.

Featured Products

ACQUITY UPC2 System <https://www.waters.com/134658367> SFC 80q/200q and 350 Preparative Systems <https://www.waters.com/10145745> Prep 100q SFC System <https://www.waters.com/10145748> 2489 UV/Visible (UV/Vis) Detector for Purification and Modular LC Systems < https://www.waters.com/134721513> 2998 Photodiode Array (PDA) Detector <https://www.waters.com/1001362> MassLynx Mass Spectrometry Software <https://www.waters.com/513164> SQ Detector 2 <https://www.waters.com/134631584> SFC Columns <https://www.waters.com/134696052>

Available for Purchase Online

Trefoil AMY1 Column, 2.5 μm, 3 mm X 150 mm, 1/pkg < https://www.waters.com/waters/partDetail.htm?partNumber=186007460> Viridis BEH Column, 130Å, 1.7 μm, 3 mm X 150 mm < https://www.waters.com/waters/partDetail.htm?locale=de_DEWaters&partNumber=186006686> Viridis BEH OBD Prep Column, 130Å, 5 μm, 19 mm X 150 mm, 1/pkg < https://www.waters.com/waters/partDetail.htm?partNumber=186005733>

720005347, April 2015

©2019 Waters Corporation. All Rights Reserved.

使用条款	I	隐私	商标	网站地图	招聘	Cookie	Cookie	设置
	沪	ICP	备06003546号-2		京公网安备 31011502007476号			