


응용 자료

Investigating Metal Containing Aggregates in Crude Oil with ACQUITY APC

Sara Gutiérrez Sama, Peter Hancock, Brice Bouyssiere, Carine Arnaudguilhem, Ryszard Lobinski, Caroline Barrere-Mangote, Pierre Giusti

Laboratory of Bioinorganic Analytical and Environmental Chemistry (LCABIE), TOTAL Refining & Chemicals, TOTAL Research & Technology, Waters Corporation, TOTAL RC – CNRS – Univ. Pau – Univ. Rouen, JOINT LABORATORY: C2MC – Complex Matrices Molecular Characterization



This is an Application Brief and does not contain a detailed Experimental section.

Abstract

In this application brief, learn how the ACQUITY APC System allows the mapping of petroleum samples faster than SEC in order to establish the structure/properties relationship, and enabling the optimization of crude oil refining.

Benefits

APC allows size profiles of undesirable metal aggregates in petroleum products to be determined in less than 10 minutes.

Introduction




Several chromatographic techniques have been developed for the fractionation of complex petroleum samples according to various physico-chemical properties, such as molecular size, the degree of aromaticity, or polarity. Gel permeation chromatography (GPC) or size exclusion chromatography (SEC) separates molecules based on their size or, more precisely, hydrodynamic volume. SEC is widely used in the crude oil industry and its use continues to increase with the new analytical challenges posed by heavy crude oil. This technique preserves the metal-ligand bond in speciation studies and also allows the study of some aggregation states. SEC has also been hyphenated to inductively coupled plasma mass spectrometry (ICP-MS) for mapping and process improvement studies, or to obtain aggregate information with specific detection of metals. A significant barrier to this technique is the lengthy analysis time (a traditional SEC analysis can take 1 to 2 hours for a single sample), as well as the solvent consumption (cost of the solvents used, cost of disposal, and associated environmental concerns).

Results and Discussion

Waters ACQUITY Advanced Polymer Chromatography (APC) System provides a breakthrough in SEC technology that defines the ultimate in size-based chromatographic separations, delivering more information about your products faster than ever before. The APC System provides unparalleled macromolecular peak

resolution, particularly for low molecular weight species, accurate and repeatable molecular weight information that is up to 5 to 20 times faster than traditional SEC methods, lower analysis costs through reduced solvent consumption and waste disposal volumes, and the ability to run diverse macromolecule applications on a single system.

Four Vacuum Residues (VR), heavy petroleum cuts with boiling point over 560 °C, were studied: VR A. from South America; VR B. from Africa; VR C. from Russia; and VR D. from Northern Europe. Separations were completed with two ACQUITY APC XT Columns of 125 Å and 450 Å (4.6 x 150 mm, 2.5 µm). 10 µL sample solutions were injected with a dilution factor of 160. The ACQUITY APC System enabled different solvents to be used and tested for sample preparation and elution without extended solvent swap out procedures. The APC RI profiles obtained are presented in Figure 1.



- Hyphenation of the APC System to ICP-HR-MS was completed for four VR samples. The size distribution profiles of V- and Ni-containing compounds showed different types of aggregates of different size.
- Significant differences of the profiles were found dependent on the origin of the sample as well as the solvent used, which seem to have an influence on the aggregation state.

Featured Products

ACQUITY Advanced Polymer Chromatography System <<https://www.waters.com/134724426>>

720005625, May 2016