

A Rapid UPLC™-MS/MS Method for the Quantification of PROTACs-3-Gefitinib and Urinary Metabolites Following Subcutaneous Administration to Male Rats

Robert S. Plumb, Nikunj Tanna

Waters Corporation

Abstract

Proteolysis Targeting Chimera (PROTAC) drugs are now well established in pharmaceutical development programs, with over 20 under clinical investigation as of the end 2023. PROTACs drugs work by mobilizing the ubiquitin–proteasome system to achieve proteasome-mediated degradation of the target protein. As such they offer a significant advantage over traditional small molecules, which inhibit protein activity, *e.g.*, complete removal of protein, reduced susceptibility to drug resistance and ability to target previously undruggable proteins. A critical part of the drug discovery development process is understanding and quantifying the in vivo fate of the molecule. As PROTACs drug molecules are essentially two active moieties, one which binds to the target protein and one which binds E3 ubiquitin ligase, connected by a linker, they are susceptible to metabolic cleavage of the linker giving rise to two active moieties which need to be detected and quantified in a DMPK study. A rapid UPLC-MS/MS method was developed and employed for the measurement of PROTACs-3-gefitinib and its major urinary metabolites following subcutaneous administration at 10 mg/kg to male rats.

Benefits

- Xevo™ TQ Absolute is a high-sensitivity MS platform with wide dynamic range for high confidence, efficient analysis, allowing for the quantification of low and high concentration analytes in one analytical batch without the need for sample dilution or reanalysis
 - UPLC Premier delivers rapid separation and quantification of PROTACs molecule and metabolites in urine
 - Reduction in non-specific binding of target analytes to transition metals in LC system, eliminating the potential for analyte loss, poor chromatographic peak shape, and irreproducible results
 - High sensitivity analysis of small sample volumes to support 3Rs (replacement, reduction, refinement) approach by facilitating serial micro sampling
-

Introduction

Proteolysis targeting chimera (PROTAC) are a relatively new class of drug molecules, first being described by Kenten *et-al* in 1999.¹ Unlike traditional small molecules, which operate by only inhibiting the enzyme function of the target protein, PROTACs produce total degradation of the entire protein, thus removing both the enzyme and non-enzyme functions. PROTACs are synthetic chemicals (small molecules) comprised of two protein binding compounds, one capable of engaging with E3 ubiquitin ligase and the other with the target protein of interest. These two molecules are covalently connected by a linker which can vary in length and chemical composition.

As part of the drug discovery and development process, it is necessary to understand the metabolism and pharmacokinetics of the compound(s) under evaluation to determine drug exposure, identify potential toxic/pharmacologically active metabolites and understand the route of elimination. Previous in vitro studies have highlighted the PROTACs linkers as being susceptible to N-dealkylation and amide hydrolysis at the points of attachment, resulting in the cleavage of the PROTACs molecule into two fragments.² Indeed, PEG based linkers were susceptible to CYP3A4 O-dealkylation offering the potential for multiple fragmentation points. These biotransformations have the potential to exhibit “on” and “off” target pharmacology or toxicity, and thus may need to be monitored and quantified, along with the dosed PROTACs, as part of any DMPK studies. As these metabolites may be present at low concentrations and have closely related structures, it is critical to have a high resolution, high sensitivity bioanalytical assay capable of separating and quantifying these metabolic products as well as the dosed compound.

Here we present a rapid, sensitive UPLC-MS/MS method using the ACQUITY™ Premier UPLC™ System and the

Xevo TQ Absolute Tandem Quadrupole Mass Spectrometer operating in positive ion ESI multiple reaction monitoring (MRM) mode for the detection and quantification of PROTAC-3-gefitinib and its metabolites in male rat urine following the subcutaneous administration to the male rat.

Experimental

Sample Description

PROTAC-3-gefitinib, Figure 1, was dosed subcutaneously at 10 mg/kg to four male rats. The dose was formulated in 5% DMSO, 40% PEG 400, 5% Cremophor, and 50% saline for injection (SFI). The rats were housed individually in a metabowl and had free access to food and water. Urine samples were collected over ice for the following time periods, 0–1 hour (n=4), 1–3 hour (n=1), 1–8 hour (n=2), 8–24 hour (n=1) following dosing. The urine samples were transferred to falcon tubes (10 mL) and stored frozen at -20 °C prior to analysis.

The study was performed under UK Home Office License PP9552589 Protocol 2-Pharmacokinetic Study, with local ethical committee clearance.

A calibration line was prepared over the range 1–1000 ng/mL by spiking PROTAC-3-gefitinib authentic standard solutions into control blank rat urine. Quality control (QC) standards were also prepared in the same manner at concentrations of 3, 75, and 800 ng/mL. Standard, QC and study samples were prepared by mixing 5 µL aliquot of urine with 20 µL cold acetonitrile: methanol (4:1) containing gefitinib D6 as an internal standard at a concentration of 50 ng/mL in 1.5 mL plastic Eppendorf tubes. The samples were vortex mixed and refrigerated at -20 °C for 1 hour. The samples were then allowed to return to room temperature and centrifuged at 14,000 g for 5 minutes. The supernatant layer was transferred to an autosampler vial for analysis.

Method Conditions

The urine extracts (1 µL) were analysed using an ACQUITY Premier UPLC System, comprising a binary solvent manager, sample manager and column manager connected to a Xevo TQ Absolute Tandem Quadrupole Mass Spectrometer. Chromatographic separations were performed using a 2.1 x 100 mm ACQUITY Premier HSS T3 1.8 µm Column maintained at 60 °C at a flow rate of 0.6 mL/min. The data was collected using MRM mode using the transitions, cone voltages and collision energies given below. Quantification was performed using gefitinib d6 stable labelled isotope as the internal standard, and 1/X weighting.

LC Conditions

LC system:	ACQUITY Premier UPLC
Vials:	LCGC Certified Clear Glass 12 x 32 mm Screw Neck Vial, Total Recovery (p/n: 186000384C)
Column(s):	ACQUITY UPLC HSS T3 Column, 100 Å, 1.8 µm, 2.1 mm x 100 mm (p/n: 186003539)
Column temperature:	60 °C
Sample temperature:	20 °C
Injection volume:	1 µL
Flow rate:	0.6 mL
Mobile phase A:	Aqueous 0.1% formic acid, 1 mM ammonium formate
Mobile phase B:	95% acetonitrile 5% water, 0.1% formic acid, 1 mM ammonium formate

Gradient Table

Time (min)	Flow rate (mL/min)	%A	%B
0	0.6	95	5
0.5	0.6	95	5
5	0.6	5	95
5.5	0.6	5	95
5.51	0.6	95	5
7	0.6	95	5

MS Conditions

MS system:	Xevo TQ Absolute
Ionization mode:	Positive ion ElectroSpray Ionization (ESI)
Capillary voltage:	1 kV
Collision energy:	34 eV
Cone voltage:	60 V

Data Management

Acquisition software:	MassLynx™ 4.2 Software
Processing software:	TargetLynx™ XS Software

PROTAC-3-Gefitinib and Metabolite MRM Transitions

Compound	Precursor ion	Product ion	Dwell time (ms)	Cone voltage	Collision energy
PROTACs	934.4	617.3	0.024	60	34
M1-EBM	521.2	87.1	0.024	60	34
M4-TDM	476.2	87.1	0.024	60	34
M5-TBM-SO ₄	697.2	87.1	0.024	60	34
M6 EBM+linker	525.2	77.0	0.024	60	34
M7 TBM-Gluc	476.1	133.1	0.024	60	34
M8-TBM-O-gluc	510.1	77.0	0.024	60	34
M9-TBM-O ₂ -Sulphate	518.1	99.1	0.024	60	34

Results and Discussion

PROTAC-3-gefitinib is a PROTAC version of gefitinib, a potent tyrosine kinase inhibitor which interrupts signalling via the epidermal growth factor receptor (EGFR) in target cells and is used to treat cancer *e.g.* non-small cell lung cancer and breast cancers. Previous studies have shown that gefitinib is well absorbed with a good bioavailability, but it is subject to extensive biotransformation in both preclinical species and humans to a large number of metabolites.^{3,4} These include O-demethylation, oxidative metabolism of the morpholine ring, and oxidative defluorination, mainly mediated via the cytochrome p450 family of enzymes.⁵ Some of these biotransformations can be further metabolized to form sulphate and glucuronide conjugates. Metabolite profiling of the urine and plasma samples obtained from dosing of PROTAC-3-gefitinib to male rats, using reversed-phase chromatography and high-resolution mass spectrometry, identified seven potentially drug related metabolites, see Table 1. These metabolites were characterized by N-dealkylation and/or hydrolysis of the aliphatic linker to yield the E3 ligase binding moiety (EBM) and the gefitinib based target binding moiety (TBM), the latter also being subject to further metabolism via conjugation to form sulphate or glucuronide metabolites. Examples of these metabolites are given in Figure 1.

Analyte	Component name	Moiety	Precursor ion	Product ion
PROTACs	PROTAC Gefitinib		934.37	617.3
M1	PROTAC Gefitinib-C ₂₂ H ₂₃ ClFN ₃ O ₃ (cleavage)+O+H ₂	EBM	521.24	87.05
M4	PROTAC Gefitinib-C ₂₅ H ₃₂ N ₄ O ₅ S(cleavage)+C ₂ H ₂ O	TBM	476.17	87.05
M5	PROTAC Gefitinib-C ₁₆ H ₁₇ N ₃ O ₂ S(cleavage)-H ₂ +SO ₃	TBM-Sulphate	697.21	87.05
M6	PROTAC Gefitinib-C ₂₂ H ₂₃ ClFN ₃ O ₄ (cleavage)+2x(-H ₂)+C ₂ H ₂ O	EBM+linker	525.21	77.04
M7	PROTAC Gefitinib-C ₃₂ H ₄₆ N ₄ O ₇ S(cleavage)+2x(-H ₂)+C ₆ H ₈ O ₆	TBM-Gluc	476.07	133.07
M8	PROTAC Gefitinib-C ₃₂ H ₄₆ N ₄ O ₆ S(cleavage)+O-H ₂ +C ₆ H ₈ O ₆	TBM-Ogluc	510.07	77.04
M9	PROTAC Gefitinib-C ₂₇ H ₃₆ N ₄ O ₅ S(cleavage)+2x(+O)+SO ₃	TBM-Sulphate	518.08	99.05

Table 1. Major plasma and urinary metabolites of PROTAC-3-gefitinib.




Figure 1. PROTACs-3-gefitinib and *in vivo* metabolites M1, M4, M5, M6, M7, M8, and M9.

PROTAC-3-Gefitinib Quantification

The urine samples were prepared by protein precipitation with acetonitrile : methanol solution (containing gefitinib stable labelled isotope (d6) (50 ng/mL) as internal standard) via vortex mixing and refrigerated at -20 °C, followed by centrifugation to yield a supernatant layer for analysis. The concentration of PROTAC-3-gefitinib

and its major metabolites were monitored using reversed-phase UPLC-MS/MS operating in +ESI MRM mode, (see method section). The MRM transitions were identified, and the cone voltages and collision energies optimized by infusion of a 100 ng/mL authentic standard using the Intellistart function within MassLynx (Waters application note, 720008229 <<https://www.waters.com/nextgen/global/library/application-notes/2024/rapid-high-sensitivity-lc-ms-ms-bioanalytical-method-for-the-simultaneous-quantification-of-gefitinib-based-protacs-3-and-gefitinib-in-rat-plasma-to-support-discovery-dmpk-studies.html>>).⁶ PROTACs-3-gefitinib eluted with retention time t_R =3.29 min, an example chromatogram for PROTACs-3-gefitinib obtained from the 8–24 hour urine sample is shown in Figure 2.




Figure 2. Example extracted ion reversed-phase +ve ESI UPLC-MS/MS chromatogram for PROTACs-3-gefitinib and internal standard from urine sample 8–24 hours.

A calibration line was constructed in urine over the concentration range 1–1000 ng/mL using a 1/x weighting and QC samples were employed at concentrations of 3, 75, and 800 ng/mL to confirm assay performance. A typical calibration line obtained from the urine analysis is shown in Figure 3. The response for the PROTACs-3-gefitinib was linear over the range of 1–1000 ng/mL with a correlation coefficient (r^2) of 0.9987 and an intercept of -0.0014. The derived calibration line was used to determine the concentration of PROTACs-3-gefitinib in the rat urine samples. The resulting mean concentration data ($n=4$) for PROTACs-3-gefitinib showed that the

elimination of the drug compound steadily increased over the period of the study and that the drug was not fully eliminated from the circulatory system by the 24 hour time point, Figure 4.

Compound name: PROTACs-3-Gefitinib
Correlation coefficient: $r = 0.999326$, $r^2 = 0.998652$
Calibration curve: $0.00358545 * x + -0.00142257$
Response type: Internal Std (Ref 1), Area * (IS Conc. / IS Area)
Curve type: Linear, Origin: Exclude, Weighting: $1/x$, Axis trans: None




Figure 3. Calibration line for PROTACs-3-gefitinib in extracted urine 1-1000 ng/mL using $1/x$ weighting.




Figure 4. Urinary elimination time course for PROTACs-3-Gefitinib following subcutaneous dose of 10 mg/kg to male rat.

Metabolite Quantification

Prior analysis of these samples using reversed-phase UPLC-HRMS showed that PROTACs-3-gefitinib was metabolized via amide hydrolysis, dealkylation of the linker chain, oxygenation of both the therapeutic binding moiety (TBM) and the E3 ligase Binding Moiety (EBM) followed by sulphate and glucuronide conjugation. A total of six major metabolites were monitored by targeted UPLC-MS/MS, in the absence of authentic standards for these metabolites the collision energy and cone voltage of PROTAC-3-gefitinib was used for the metabolites. The drug related metabolites eluted between $t_R=1.94$ and 3.27 minutes, specifically M1 $t_R=2.74$, M4 $t_R=1.94$, M6 $t_R=3.27$, M7 $t_R=2.86$, M8 $t_R=2.40$, and M9 $t_R=2.41$. Example extracted positive ion chromatograms for the metabolites; cleavage of EBM M4, (476.16 \rightarrow 89.06); cleavage of the gefitinib TBM and oxygenation of the EBM M1, (531.24 \rightarrow 89.06); cleavage of gefitinib TBM and linker M6 (525.2 \rightarrow 77.04) and cleavage of the EBM followed by glucuronidation of the TBM M7, are given in Figures 5–8. The UPLC-MS/MS data for these metabolites

showed that the system produced excellent peak shape and good resolution of the drug metabolites from the dosed compound and endogenous urinary components. In the absence of authentic standards, the drug metabolites were quantified against the PROTACs-3-gefitinib calibration line. The high sensitivity and wide dynamic range of the Xevo TQ Absolute Tandem Quadrupole Mass Spectrometer allowed for the quantification of both the low concentration PROTACs-3-gefitinib and the low concentration metabolites (M8, M6) as well as the higher concentration metabolites (M9, M4, M1, and M7) in a single analytical batch.




Figure 5. Example extracted ion chromatogram (476.16 → 89.06) for M4 metabolite and internal standard from urine sample 8-24 hours.




Figure 6. Example extracted ion chromatogram (531.24 → 89.06) for M1 metabolite and internal standard from urine sample 8–24 hours.




Figure 7. Example extracted ion chromatogram (525.2 → 77.04) for M6 metabolite from urine sample 1-8 hours.




Figure 8. Example extracted ion chromatogram (467.07 → 133.07) for M7 metabolite and internal standard from urine sample 1–8 hours.

The resulting metabolite mass elimination data (concentration x volume of urine) is summarized in Figure 9. As can be seen from this data the metabolites M9, M4, M1, and M7 were responsible for the largest proportion of the drug related eliminated material on each sampling occasions. The total mass eliminated for the metabolites M9, M4, M1, and M7 over the 24 h sample period was 275.2, 216.5, 223.3, and 54.2 µg, respectively. The metabolites M9, M7 are the sulphate and glucuronide conjugate metabolites of the TBM following dealkylation of the linker, whilst the metabolites M1, M4 are the TBM, and EBM products resulting from the from the cleavage of the PROTACs linker (O-dealkylation). It is interesting to note that the two cleavage metabolites (M1, M4) are likely formed by the same metabolic cleavage reaction. The metabolites M8, M6, and PROTACs-3-gefitinib were present at significantly lower concentrations and contributed much less to the mass of eliminated material over the 24 hour time period.




Figure 9. Urinary elimination (μg) of PROTACs-3-gefitinib and its metabolites following subcutaneous administration at 10 mg/kg to male rats.

Conclusion

PROteolysis Targeting Chimeras are a new class of drug molecules which offer great promise to overcome the disadvantages of typical small molecule drugs, such as drug resistance, toxicity, and inability to address all the druggable proteome, whilst retaining the advantages of simple chemical synthesis, easy scale up and patient friendly administration. In this study we have employed sub 2 μm reversed-phase chromatography coupled to tandem quadrupole MS operated in multiple reaction monitoring mode to quantify PROTAC-3-gefitinib and its metabolites in urine following subcutaneous administration at 10 mg/kg to the male rat. The high sensitivity and wide linear dynamic range of the Xevo TQ Absolute facilitated the quantification of both the low concentration PROTAC-3-gefitinib drug and the M6 and M8 metabolites as well as the higher concentration M1, M4, M7, and M9 metabolites in one single batch. The high-resolution separation offered by the ACQUITY Premier UPLC System allowed for the resolution of PROTAC-3-gefitinib from the metabolites as well as the metabolites from each other, thus improving the specificity of the quantification. The resulting data showed that the majority of the

drug related material eliminated in the urine was in the form of O-glucuronide, sulfate, and O-dealkylation cleavage metabolites. The data also showed that the drug related material had not been completely eliminated from the system by the 24 hour sampling time point.

References

1. Kenten J.H, Roberts S.F. Controlling Protein Levels in Eucaryotic Organisms, US Patent US6306663B1.
2. Goracci L, Desantis J, Valeri A, Castellani B, Eleuteri M, Cruciani G. Understanding the Metabolism of Proteolysis Targeting Chimeras (PROTACs): The Next Step Toward Pharmaceutical Applications. *J Med Chem* . 2020 22;63(20):11615–11638. doi: 10.1021/acs.jmedchem.0c00793.
3. Molloy B.J., King A., Mullin L., Gethings L.A., Riley R., Plumb R., Wilson I.D. Rapid Determination of the Pharmacokinetics and Metabolic Fate of Gefitinib in the Mouse Using a Combination of UPLC/MS/MS, UPLC/QToF/MS, and Ion Mobility (IM)-Enabled UPLC/QToF/MS. *Xenobiotica*. 2021;51:434–446. doi: 10.1080/00498254.2020.1859643.
4. McKillop D., Hutchison M., Partridge E.A., Bushby N., Cooper C.M.F., Clarkson-Jones J.A., Herron W., Swaisland H.C. Metabolic Disposition of gefitinib, an Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor, in Rat, Dog, and Man. *Xenobiotica*. 2004;34:917–934. doi: 10.1080/00498250400009171.
5. Liu X., Lu Y., Guan X., Dong B., Chavan H., Wang J., Zhang Y., Krishnamurthy P., Li F. Metabolomics Reveals the Formation of Aldehydes and Iminium in Gefitinib Metabolism. *Biochem. Pharmacol*. 2015;97:111–121. doi: 10.1016/j.bcp.2015.07.010.
6. Rapid High Sensitivity LC-MS/MS Bioanalytical Method for the Simultaneous Quantification of Gefitinib Based PROTACs-3 and Gefitinib in Rat Plasma to Support Discovery DMPK Studies, Waters Application Note. 720008229 <<https://www.waters.com/nextgen/global/library/application-notes/2024/rapid-high-sensitivity-lc-ms-ms-bioanalytical-method-for-the-simultaneous-quantification-of-gefitinib-based-protacs-3-and-gefitinib-in-rat-plasma-to-support-discovery-dmpk-studies.html>> . January 2024.

Featured Products

[ACQUITY Premier System <](#)

<https://www.waters.com/nextgen/global/products/chromatography/chromatography-systems/acquity-premier-system.html>>

[Xevo TQ Absolute Triple Quadrupole Mass Spectrometry <](#)

<https://www.waters.com/nextgen/global/products/mass-spectrometry/mass-spectrometry-systems/xevo-tq-absolute.html>>

[MassLynx Mass Spectrometry Software <https://www.waters.com/nextgen/global/products/informatics-and-software/mass-spectrometry-software/masslynx-mass-spectrometry-software.html>](#)

[TargetLynx <https://www.waters.com/513791>](#)

720008484, October 2024



© 2024 Waters Corporation. All Rights Reserved.

[使用條款](#) [隱私權政策](#) [商標](#) [就業機會](#) [法律和隱私權聲明](#) [Cookie](#) [Cookie偏好設定](#)